

NUTRITION INFORMATION

ZeroTP Infant Mix LCP

		100 g powder	100 ml ready-to-use*
• Energy	kJ	2123	318
	kcal	508	76
• Fat	g	28	4
of which	saturates	5	0,8
mono-unsaturates	g	11	1,6
polyunsaturates	g	12	1,7
• Carbohydrate	g	54	8
of which	sugars	25	4
	starch	8	1,2
• Protein equiv.	g	11	1,7
Amino acids	g	13	2
• Salt	g	0,6	0,1

FURTHER NUTRITION INFORMATION

• Lactose	g	24	3,5
• Maltodextrin	g	22	3

- Standard dilution: 15 g ZeroTP Infant Mix LCP + 90 ml drinking water = 100 ml ready-to-use formula
- Osmolality: 350 - 360 mosmol/ kg drinking water
- The enclosed scoop measures approx. 3,7 g powder.



ZeroTP Infant Mix LCP

- ✓ 1 x 500 g tin
- ✓ 6 x 500 g tin

✓ ZeroTP Infant Mix LCP

Soon attractive also for your child!

Tastes good!

Large choice of protein-reduced and low protein foodstuffs

www.metax-shop.org

hotline 008000-9963829
(free of charge from A, D, NL)
tel +49 (0) 8432 - 94 86 0
fax +49 (0) 8432 - 94 86 19
eMail info@metax.org
visit us on www.metax.org

metaX Institut fuer Diabetik GmbH
Am Strassbach 5 · 61169 Friedberg GERMANY

16/07/08 d 14:04:09

INFORMATION FOR PARENTS



ZeroTP Infant Mix LCP

- food for special medical purposes
- free of Tyrosine and Phenylalanine (TP)
- for the dietary management of Tyrosinemia

formula for Infants

- highly purified L-amino acids
- with vitamins, minerals and trace elements in adequate amounts
- with LCPUFA-lipids
- with lactose, according to mother's milk standard

✓ Definition and Indication

- **ZeroTP Infant Mix LCP** is an infant formula free of Tyrosine and Phenylalanine (TP) for use in the dietary management of Tyrosinemia in infants. **ZeroTP Infant Mix LCP** consists of amino acids, fat and carbohydrate and is enriched with vitamins, minerals and trace elements. Like breast milk **ZeroTP Infant Mix LCP** contains lactose as main carbohydrate and is enriched with LCPUFA-lipids.

✓ Dosage

- The correct dosage of **ZeroTP Infant Mix LCP** is determined by the physician and/ or dietitian and should be re-examined and adjusted in accordance with the results of regular monitoring. The dosage depends on age, body weight, protein requirements and the infant's individual metabolic situation. The daily amount is determined individually and is given to the infant in combination with calculated amounts of breast milk and/ or regular infant formula and/ or low protein weaning food (beikost).

✓ Directions for use

- The total daily quantities of **ZeroTP Infant Mix LCP** and other foods, such as breast milk, infant formula or low protein weaning food, to be used in the dietary management of Tyrosinemia, are determined in a diet plan. The daily amount of **ZeroTP Infant Mix LCP** should be given in 3 - 5 portions distributed during the day, as a bottle feed or pap.
- When prepared with drinking water, **ZeroTP Infant Mix LCP** gives a bottle feed free of Tyrosine and Phenylalanine.
- **ZeroTP Infant Mix LCP** must not be given without other calculated, protein containing foods (e.g. breast milk or infant formula) over a longer period of time. Medical supervision is absolutely essential.

✓ Preparation



- Clean and boil glass bottle and teat before use.
- Boil the drinking water and allow to cool down to 40 °C.
- Pour required amount of boiled water into the bottle.
- Level the scoop with the back of a knife. Only use the scoop enclosed in the tin!
- Put the lid safely on the bottle and shake.
- Open the bottle and fix the teat. Check drinking temperature (cheek test).
- **At best always weigh out the amount needed for the preparation of the bottle feed with ZeroTP Infant Mix LCP!**
- Always prepare the bottle feed or the pap with **ZeroTP Infant Mix LCP** freshly. Do not reuse any leftovers.
- For preparing a pap, **ZeroTP Infant Mix LCP** can be mixed additionally with calculated amounts of starch, low protein biscuits and/ or fruits.

Advice for dental care

Carbohydrates are indispensable for the nutrition of a child. Like mother's milk, **ZeroTP Infant Mix LCP** contains carbohydrates. Incorrect use of the bottle (frequent or continuous sucking) can result in tooth decay (caries) and related damages to health. Thus a bottle feed should only last for the time required for the food intake. Do not leave the bottle with your child for continuous sucking.

✓ ZeroTP Infant Mix LCP

✓ ZeroTP Infant Mix LCP

✓ ZeroTP Infant Mix LCP